

The information in this guide is intended only for patients who have been prescribed OVIDREL® pre-filled syringe by a doctor experienced in treating infertility. Before injecting with the OVIDREL® pre-filled syringe, please read the leaflet inside the OVIDREL® syringe box.


# OVIDREL®

choriogonadotropin alfa (rch)

Pre-filled Syringe

## Administration & storage instructions

One pre-filled syringe  
of OVIDREL® 250 µg




Alcohol wipes  
(not included in the package)

**ADMINISTRATION – REVIEW THESE  
STEPS BEFORE YOU ADMINISTER OVIDREL®**

### Before you begin:

- Wash your hands well with soap & water
- Lay out the above items on a clean surface



### STEP 1

Your doctor, nurse or pharmacist will have already advised you where to inject, e.g. tummy or front of thigh, as shown in the diagram to the left.



### STEP 6

Once the needle is inserted into the tissue all the way, inject the solution by pushing gently on the plunger. Take as much time as you need to inject all the solution.



### STEP 2

Wipe the chosen area with an alcohol swab.



### STEP 7

Withdraw the needle and clean the site with an alcohol swab using a circular motion.

Once you have finished your injection, immediately discard the empty syringes in a sharps container. Any unused solution must be discarded.



### STEP 3

Pick up the syringe and remove the cap from the needle. (Please note the syringe may appear differently to the syringe pictured on the left).

**Before injecting with the OVIDREL<sup>®</sup> pre-filled syringe, please read the leaflet inside the OVIDREL<sup>®</sup> syringe box.**



### STEP 4


Invert the needle and hold as if "throwing a dart".

With your other hand, firmly pinch the skin together to make a little elevation at the injection site.

### Reporting of side effects:

If you get any side effects, talk to your doctor, pharmacist or nurse and report the side effects to MERCK at 1800 633 463 or by emailing [ICSR\\_ANZ@merckgroup.com](mailto:ICSR_ANZ@merckgroup.com).

You can also report side effects directly to TGA at <https://www.tga.gov.au/reporting-problems>.



### STEP 5

Using a "dart-like motion", insert the needle at a 45° to 90° angle.

**IMPORTANT: To inject, you need very little force but quick action.**